



Modeling the Atmospheric Transport and Deposition of Mercury to the Great Lakes

**Dr. Mark Cohen
NOAA Air Resources Laboratory (ARL)
College Park, MD, USA**

**NOAA Air Resources Laboratory Seminar
February 7, 2013, NCWCP, College Park MD**



Acknowledgements

**HYSPLIT
Development
Team at ARL:**

**Roland Draxler
Glenn Rolph
Barbara Stunder
Ariel Stein
Fantine Ngan**

**Mercury
Research
Team at ARL:**

**Winston Luke
Paul Kelley
Steve Brooks
Xinrong Ren**

**IT Team
at ARL:**

**Rick Jiang
Yan Huang**


Funding:

**Great Lakes Restoration Initiative,
via Interagency Agreement with USEPA**

*+ numerous collaborations with
external partners involving:*

- emissions inventory data for model input, and
- atmospheric measurement data for model evaluation

Mercury in the Environment




Biodiversity Research Institute (2011):

http://www.briloon.org/uploads/hgconnections/glmc/GLMC_FinalReport.pdf



**0.05 ppm level
recommended by the
Great Lakes Fish Advisory
Workgroup (2007)**


Mercury in Great Lakes Fish



Evers, D.C., et al. (2011). *Great Lakes Mercury Connections: The Extent and Effects of Mercury Pollution in the Great Lakes Region*. Biodiversity Research Institute. Gorham, Maine. Report BRI 2011-18. 44 pages.



Temporal trends of mercury in Lake Erie 45–55 cm walleye collected between 1990–2007 (Bhavsar et al., Environ. Sci. Technol. 2010, 44, 3273-3279)



Atmospheric deposition is believed to be the largest current mercury loading pathway to the Great Lakes...

➤ **How much is deposited and where does it come from?**

(...this information can *only* be obtained via modeling...)



Different “forms” of mercury in the atmosphere

Elemental Mercury -- Hg(0)


- most of total Hg in atmosphere
- doesn't easily dry or wet deposit
- globally distributed

Reactive Gaseous Mercury -- RGM

- a few % of total atmos. Hg
- oxidized Hg (HgCl_2 , others)
- *very* water soluble and “sticky”
- bioavailable


Particulate Mercury -- Hg(p)

- a few % of total atmos. Hg
- Hg in/on atmos. particles
- atmos. lifetime 1~ 2 weeks
- bioavailability?





Starting point: where is mercury emitted to the air?


2005 Atmospheric Mercury Emissions from Large Point Sources



2005 Atmospheric Mercury Emissions (Direct Anthropogenic + Re-emit + Natural)





**Atmospheric mercury emissions (kg/yr)
from all sources in each 2x2 degree grid cell**



Next step: What happens to the mercury after it is emitted?

HYSPLIT-Hg Lagrangian Puff Atmospheric Fate and Transport Model







When puffs grow to sizes large relative to the meteorological data grid, they split, horizontally and/or vertically

To simulate the global transport of mercury, puffs are transferred Eulerian grid after a specified time downwind (~2 weeks), and the mercury is simulated that grid from then on


This is how we model the local & **regional** impacts.

But for **global** modeling, puff splitting overwhelms computational resources




● Deposition explicitly modeled to actual lake/watershed areas

- As opposed to the usual practice of ascribing portions of gridded deposition to these areas in a post-processing step



Illustrative simulation of reactive gaseous mercury (RGM) emissions from one power plant on the shore of Lake Erie: hourly deposition estimates for the first two weeks in May 2005

Deposition (ng/m²-hr) at ground-level
Integrated from 0000 01 May to 0100 01 May 05 (UTC)
RGM Release started at 0000 01 May 05 (UTC)







Computational Challenge

- ➔ **350,000 sources in global emissions inventory**
- ➔ **Would like to keep track of each source individually**
- ➔ **typical one-year simulation takes
~96 processor hours**
- ➔ **~3800 processor years, if ran explicit
simulation for each source**
- ➔ **~240 years on 16-processor workstation**

Spatial Interpolation



Chemical Interpolation




Impact of Source
Emitting
30% Hg(0)
50% Hg(II)
20% Hg(p)

$$\begin{aligned}
 &= 0.3 \times \text{Impact of Source Emitting Pure Hg(0)} \\
 &\quad + \\
 &0.5 \times \text{Impact of Source Emitting Pure Hg(II)} \\
 &\quad + \\
 &0.2 \times \text{Impact of Source Emitting Pure Hg(p)}
 \end{aligned}$$


Standard Points in North America

For each standard source location, we do three unit-emissions simulations:

- *pure Hg(0),*
- *pure HgII (RGM)*
- *pure Hg(p)*




Atmospheric mercury emissions (kg/yr) from all sources in each 1x1 deg grid cell



Standard Points Outside of North America

...three unit-emissions simulations for each location






Computational Solution


- ➔ **This analysis done with 136 standard source locations**
- ➔ **3 unit emissions simulations from each location (Hg(0), RGM, and Hg(p))**
- ➔ **~4.5 processor years**
- ➔ **~3.5 months on 16-processor workstation**

instead of 240 years ... almost 1000x less!


an example for one source... the Monroe coal-fired power plant on the shore of Lake Erie



- Monroe emitted 561 kg of mercury in 2005 (EPA's National Emissions Inventory)
- How much of this mercury was deposited into Lake Erie and its watershed?



- Monroe emitted 561 kg of mercury in 2005 (EPA's National Emissions Inventory)
- Modeling results for this specific source:
 - 24 kg (~4%) of this emitted mercury was deposited directly into Lake Erie
 - 107 kg (~19%) of this emitted mercury was deposited in the Lake Erie Watershed
- We make this same type of estimate for *every source* in the national and global emissions inventories used as model input... *using spatial and chemical interpolation*






Outline of Modeling Analysis


- **HYSPLIT**
 - **HYSPLIT-Hg** (with mercury-specific chemistry, ...)
 - **Unit Emissions Simulations** of Hg(0), Hg(II) and Hg(p) from an array of standard source locations
 - **Emissions Inventory** – emissions of Hg(0), Hg(II), and Hg(p) from sources at specified latitudes and longitudes
 - **“Multiplication”** of emissions inventory by array of unit emissions simulations using spatial and chemical interpolation
-
- **Evaluate** overall model results: compare against ambient measurements
 - **Source-attribution** results for deposition to selected receptors

After all the standard source simulations have been run, and the impacts of each of the ~350,000 sources worldwide are estimated using spatial and chemical interpolation, is the model giving reasonable results?


Modeled vs. Measured Wet Deposition of Mercury at Sites in the Great Lakes Region



Standard source locations, MDN sites, and mercury emissions in the Great Lakes region



Atmospheric mercury emissions (kg/yr) from all sources in each 1x1 deg grid cell



● Standard source locations for HYSPLIT-Hg simulations

□ MDN sites in the "western" Great Lakes region



■ MDN sites in the "eastern" Great Lakes region




Outline of Modeling Analysis

- **HYSPLIT**
 - **HYSPLIT-Hg** (with mercury-specific chemistry, ...)
 - **Unit Emissions Simulations** of Hg(0), Hg(II) and Hg(p) from an array of standard source locations
 - **Emissions Inventory** – emissions of Hg(0), Hg(II), and Hg(p) from sources at specified latitudes and longitudes
 - **“Multiplication”** of emissions inventory by array of unit emissions simulations using spatial and chemical interpolation
 - **Evaluate** overall model results: compare against ambient measurements
-
- **Source-attribution** results for deposition to selected receptors


2005 Atmospheric Mercury Emissions (Direct Anthropogenic + Re-emit + Natural)



**Atmospheric mercury emissions (kg/yr)
from all sources in each 2x2 degree grid cell**




Keep track of the contributions from each source, and add them up




Geographical Distribution of 2005 Atmospheric Mercury Deposition Contributions to Lake Erie

A tiny fraction of 2005 global mercury emissions within 500 km of Lake Erie




Modeling results show that these “regional” emissions are responsible for a large fraction of the modeled 2005 atmospheric deposition



Important policy implications!




**Cumulative
Fraction
of Total
Modeled
Mercury
Deposition
to Lake Erie
(2005)**




Top 50 Atmospheric Deposition Contributors to Lake Erie

Cumulative Fraction of Total Modeled Deposition (2005)



Rank of Source's Atmospheric Mercury Deposition Contribution to Lake Erie


Sources of Mercury Deposition to the Great Lakes Basin 2005 Baseline Analysis



Total = 11,300 kg/yr




Sources of Mercury Deposition to the Lake Erie Basin 2005 Baseline Analysis




Total = 2,300 kg/yr

Sources of Mercury Deposition to the Great Lakes Basin 2005 Baseline Analysis



Total = 11,300 kg/yr

Sources of Mercury Deposition to the Lake Erie Basin 2005 Baseline Analysis




Total = 2,300 kg/yr



A multi-phase project

ARL's GLRI Atmospheric Mercury Modeling Project





Phase 1: Baseline Analysis for 2005

(Final Report Completed December 2011)

- 2005 was chosen as the analysis year, because 2005 was the latest year for which comprehensive mercury emissions inventory data were available at the start of this project
- Using 2005 meteorological data and emissions, the deposition and source-attribution for this deposition to each Great Lake and its watershed was estimated
- The model results were ground-truthed against 2005 Mercury Deposition Network data from sites in the Great Lakes region



Modeling Atmospheric Mercury Deposition to the Great Lakes.

Final Report for work conducted with FY2010 funding from the Great Lakes Restoration Initiative. December 16, 2011.

Mark Cohen, Roland Draxler, Richard Artz. NOAA Air Resources Laboratory, Silver Spring, MD, USA. 160 pages.

http://www.arl.noaa.gov/documents/reports/GLRI_FY2010_Atmospheric_Mercury_Final_Report_2011_Dec_16.pdf

http://www.arl.noaa.gov/documents/reports/Figures_Tables_GLRI_NOAA_Atmos_Mercury_Report_Dec_16_2011.pptx

One-page summary:

http://www.arl.noaa.gov/documents/reports/GLRI_Atmos_Mercury_Summary.pdf



Some Key Features of this Analysis

- **Deposition explicitly modeled to actual lake/watershed areas**
 - As opposed to the usual practice of ascribing portions of gridded deposition to these areas in a post-processing step

- **Combination of Lagrangian & Eulerian modeling**
 - allows accurate and computationally efficient estimates of the fate and transport of atmospheric mercury over all relevant length scales – from “local” to global.

- **Uniquely detailed source-attribution information is created**
 - deposition contribution to each Great Lakes and watersheds from each source in the emissions inventories used is estimated individually
 - The level of source discrimination is only limited by the detail in the emissions inventories
 - Source-type breakdowns not possible in this 1st phase for global sources, because the global emissions inventory available did not have source-type breakdowns for each grid square



Some Key Findings of this Analysis

- **“Single Source” results illustrate source-receptor relationships**
 - For example, a “typical” coal-fired power plant near Lake Erie may contribute on the order of 1000x the mercury – for the same emissions – as a comparable facility in China.

- **Regional, national, & global mercury emissions are all important contributors to mercury deposition in the Great Lakes Basin**
 - For Lakes Erie and Ontario, the U.S. contribution is at its most significant
 - For Lakes Huron and Superior, the U.S. contribution is less significant.
 - Local & regional sources have a much greater atmospheric deposition contributions than their emissions, as a fraction of total global mercury emissions, would suggest.



Some Key Findings of this Analysis (...continued)

● Reasonable agreement with measurements

- Despite numerous uncertainties in model input data and other modeling aspects
- Comparison at sites where significant computational resources were expended – corresponding to regions that were the most important for estimating deposition to the Great Lakes and their watersheds – showed good consistency between model predictions and measured quantities.
- For a smaller subset of sites generally downwind of the Great Lakes (in regions not expected to contribute most significantly to Great Lakes atmospheric deposition), less computational resources were expended, and the comparison showed moderate, but understandable, discrepancies.




Phase 2: Sensitivity Analysis + Extended Model Evaluation **(current work, with GLRI FY11 funding)**

- **Examining the influence of uncertainties on the modeling results, by varying critical model parameters, algorithms, and inputs, and analyzing the resulting differences in results**
- **Ground-truthing the model against additional ambient monitoring data, e.g., ambient mercury air concentration measurements and wet deposition data not included in the Mercury Deposition Network (MDN)**


Fraction of RGM emissions deposited in Lake Erie from a hypothetical source in Detroit

Release height
50 m vs 250 m

Met data
(EDAS vs. NARR)



Fraction
of Hg(0)
emissions
deposited in
Lake Erie
from a
hypothetical
source
in China





Phase 3: Scenarios

(next year's work, with GLRI FY12 funding)

- A modeling analyses such as this is the *only* way to quantitatively examine the potential consequences of alternative future emissions scenarios
- We will work with EPA and other Great Lakes Stakeholders to identify and specify the most policy relevant scenarios to examine
- For each scenario, we will estimate the amount of atmospheric deposition to each of the Great Lakes and their watersheds, along with the detailed source-attribution for this deposition


Thanks!






EXTRA SLIDES

Atmospheric Mercury Deposition ($\mu\text{g}/\text{m}^2\text{-yr}$)







(Evolving) Atmospheric Chemical Reaction Scheme for Mercury

Reaction	Rate	Units	Reference
GAS PHASE REACTIONS			
? $\text{Hg}^0 + \text{O}_3 \rightarrow \text{Hg(p)}$	3.0E-20	cm ³ /molec-sec	Hall (1995)
$\text{Hg}^0 + \text{HCl} \rightarrow \text{HgCl}_2$	1.0E-19	cm ³ /molec-sec	Hall and Bloom (1993)
$\text{Hg}^0 + \text{H}_2\text{O}_2 \rightarrow \text{Hg(p)}$	8.5E-19	cm ³ /molec-sec	Tokos et al. (1998) (upper limit based on experiments)
$\text{Hg}^0 + \text{Cl}_2 \rightarrow \text{HgCl}_2$	4.0E-18	cm ³ /molec-sec	Calhoun and Prestbo (2001)
? $\text{Hg}^0 + \text{OH} \rightarrow \text{Hg(p)}$	8.7E-14	cm ³ /molec-sec	Sommar et al. (2001)
new $\text{Hg}^0 + \text{Br} \rightarrow \text{HgBr}_2$			
AQUEOUS PHASE REACTIONS			
$\text{Hg}^0 + \text{O}_3 \rightarrow \text{Hg}^{+2}$	4.7E+7	(molar-sec) ⁻¹	Munthe (1992)
$\text{Hg}^0 + \text{OH} \rightarrow \text{Hg}^{+2}$	2.0E+9	(molar-sec) ⁻¹	Lin and Pehkonen(1997)
$\text{HgSO}_3 \rightarrow \text{Hg}^0$	$T^*e^{((31.971*T)-12595.0)/T}$ sec ⁻¹ [T = temperature (K)]		Van Loon et al. (2002)
? $\text{Hg(II)} + \text{HO}_2 \rightarrow \text{Hg}^0$	~ 0	(molar-sec) ⁻¹	Gardfeldt & Jonnson (2003)
$\text{Hg}^0 + \text{HOCl} \rightarrow \text{Hg}^{+2}$	2.1E+6	(molar-sec) ⁻¹	Lin and Pehkonen(1998)
$\text{Hg}^0 + \text{OCl}^{-1} \rightarrow \text{Hg}^{+2}$	2.0E+6	(molar-sec) ⁻¹	Lin and Pehkonen(1998)
$\text{Hg(II)} \leftrightarrow \text{Hg(II)}_{(\text{soot})}$	9.0E+2	liters/gram; t = 1/hour	eqlbm: Seigneur et al. (1998) rate: Bullock & Brehme (2002).
$\text{Hg}^{+2} + \text{h}\nu \rightarrow \text{Hg}^0$	6.0E-7	(sec) ⁻¹ (maximum)	Xiao et al. (1994); Bullock and Brehme (2002)



What year to model?

● Mercury Emissions Inventory

- U.S. anthropogenic emissions inventory
- Canadian anthropogenic emissions inventory
- Mexican anthropogenic emissions inventory
- Global anthropogenic emissions inventory
- Natural emissions inventory
- Re-emissions inventory

● Ambient Data for Model Evaluation

- Wet deposition (Mercury Deposition Network)
- “Speciated” Air Concentrations

● Meteorological Data to drive model


- NCEP/NCAR Global Reanalysis (2.5 deg)
- NCEP EDAS 40km North American Domain
- North American Regional Reanalysis (NARR)

**Dataset
Available
for 2005**

**Need all
of these
datasets
for the
same year**


***2005 chosen
for baseline
analysis***

Over the entire modeling period
(e.g., one year), puffs are released
at periodic intervals
(e.g., once every 7 hours).





Getting good ground-truthing results harder than estimating deposition to the Great Lakes




One Standard Source Location (green dot) would do a decent job of estimating deposition to the receptor, for all of the hypothetical, "actual" source locations shown (numbered boxes)

But the same Standard Source Location would be completely inadequate to estimate deposition and concentrations at the monitoring site (red star)


Standard Source Locations for Illustrative Modeling Results

Standard Source Locations for which Illustrative Modeling Results will be Shown





Lake Erie Transfer Flux Coefficients for two kinds of Generic Coal-Fired Power Plants (logarithmic scale)




The "Transfer Flux Coefficient" is calculated as the atmospheric deposition flux to a given receptor (in this case, Lake Erie) in units of g/km²-yr, divided by the total emissions from the source, in units of g/yr.

With this transfer flux coefficient, if one knows the emissions of the source in the given location, then the atmospheric deposition flux impact of the source on the receptor can be estimated, by simply multiplying the emissions by the transfer flux coefficient.



Lake Erie Transfer Flux Coefficients for two kinds of Generic Coal-Fired Power Plants (linear scale)



The "Transfer Flux Coefficient" is calculated as the atmospheric deposition flux to a given receptor (in this case, Lake Erie) in units of $\text{g}/\text{km}^2\text{-yr}$, divided by the total emissions from the source, in units of g/yr .

With this transfer flux coefficient, if one knows the emissions of the source in the given location, then the atmospheric deposition flux impact of the source on the receptor can be estimated, by simply multiplying the emissions by the transfer flux coefficient.



In order to conveniently compare different model results,
a “transfer flux coefficient” X will be used,
defined as the following:

$$X = \frac{\text{deposition flux rate}}{\text{emissions rate}} = \frac{\frac{\text{grams Hg deposited per year}}{\text{km}^2 \text{ of receptor area}}}{\text{grams Hg emitted per year from the source}} [=] \frac{1}{\text{km}^2}$$

$$\text{deposition flux rate} = \frac{\text{grams Hg deposited per year}}{\text{km}^2 \text{ of receptor area}} \left(\frac{\text{g}}{\text{km}^2 \text{ yr}} \right)$$

$$= \text{transfer flux coefficient} \left(\frac{1}{\text{km}^2} \right) * \text{source mercury emissions} \left(\frac{\text{g}}{\text{yr}} \right)$$




$$\text{deposition flux rate} = \frac{\text{grams Hg deposited per year}}{\text{km}^2 \text{ of receptor area}} \left(\frac{\text{g}}{\text{km}^2 \text{ yr}} \right)$$

$$= \text{transfer flux coefficient} \left(\frac{1}{\text{km}^2} \right) * \text{source mercury emissions} \left(\frac{\text{g}}{\text{yr}} \right)$$

Transfer Flux Coefficients For Pure Elemental Mercury Emissions at an Illustrative Subset of Standard Source Locations, for Deposition Flux Contributions to Lake Erie

Transfer Flux Coefficient "X" for Elemental Mercury Emissions from Selected Locations to Lake Erie


$$X = \frac{(\text{grams Hg deposited per year}) / (\text{km}^2)}{(\text{grams Hg emitted per year from the source})} [=] 1/\text{km}^2$$




Transfer Flux Coefficients For Pure Reactive Gaseous Mercury Emissions at an Illustrative Subset of Standard Source Locations, for Deposition Flux Contributions to Lake Erie

Transfer Flux Coeff. "X" for Reactive Gaseous Mercury Emissions from Selected Locations to Lake Erie

$$X = \frac{(\text{grams Hg deposited per year}) / (\text{km}^2)}{(\text{grams Hg emitted per year from the source})} [=] 1/\text{km}^2$$



Transfer Flux Coefficients For Hg(0), Hg(II), and Hg(p) to Lake Erie (logarithmic scale)




The "Transfer Flux Coefficient" is calculated as the atmospheric deposition flux to a given receptor (in this case, Lake Erie) in units of g/km²-yr, divided by the total emissions from the source, in units of g/yr.

With this transfer flux coefficient, if one knows the emissions of the source in the given location, then the atmospheric deposition flux impact of the source on the receptor can be estimated, by simply multiplying the emissions by the transfer flux coefficient.





Transfer Flux Coefficients For Hg(0), Hg(II), and Hg(p) to Lake Erie (linear scale)




The "Transfer Flux Coefficient" is calculated as the atmospheric deposition flux to a given receptor (in this case, Lake Erie) in units of g/km2-yr, divided by the total emissions from the source, in units of g/yr.

With this transfer flux coefficient, if one knows the emissions of the source in the given location, then the atmospheric deposition flux impact of the source on the receptor can be estimated, by simply multiplying the emissions by the transfer flux coefficient.



Anthropogenic Mercury Emissions (ca. 2005)



Atmospheric mercury emissions (kg/yr) from direct anthropogenic sources in each 2x2 degree grid cell



Natural mercury emissions



**Atmospheric mercury emissions (kg/yr)
from natural sources in each 2x2 degree grid cell**




Figure 55. Mercury Deposition Network Sites in the Great Lakes Region Considered in an Initial Model Evaluation Analysis






Figure 56. Comparison of Total 2005 Precipitation Measured at each of the Great-Lakes Region MDN Sites with the Precipitation in the Meteorological Datasets Used as Inputs to this Modeling Study






Comparison of 2005 precipitation total as measured at MDN sites in the Great Lakes region (circles) with precipitation totals assembled by the PRISM Climate Group, Oregon State University




Modeled vs. Measured Wet Deposition of Mercury at Sites in the Great Lakes Region



Modeled vs. Measured Wet Deposition of Mercury at Sites in the Great Lakes Region

2005 Total Wet Mercury Deposition
($\mu\text{g}/\text{m}^2\text{-yr}$)







Summary of Mercury Emissions Inventories Used in GLRI Analysis

Inventory	domain	Number of records	Hg(0) emissions (Mg/yr)	RGM emissions (Mg/yr)	Hg(p) emissions (Mg/yr)	Total mercury emissions (Mg/yr)
U.S. Point Sources	United States	19,353	50.6	35.5	9.1	95
U.S. Area Sources	United States	44,848	4.5	1.8	1.1	7.4
Canadian Point Sources	Canada	166	3.0	1.7	0.4	5.1
Canadian Area Sources	Canada	12,372	1.0	0.96	0.42	2.4
Mexican Point Sources	Mexico	268	28	0.81	0.46	29
Mexican Area Sources	Mexico	160	1.25	0.38	0.25	1.9
Global Anthropogenic Sources not in U.S., Canada, or Mexico	Global, except for the U.S., Canada, and Mexico	52,173	1,239	434	113	1,786
Global Re-emissions from Land	Global land (and freshwater) surfaces	129,180	750	0	0	750
Global Re-emissions from the Ocean	Global oceans	43,324	1,250	0	0	1,250
Global Natural Sources	Global	64,800	1,800	0	0	1,800
Total		366,804	5,127	475	125	5,728



Model-estimated 2005 deposition to the Great Lakes Basin from countries with the highest modeled contribution from direct and re-emitted anthropogenic sources





Model-estimated *per capita* 2005 deposition to the Great Lakes Basin from countries with the highest modeled contribution from direct & re-emitted anthropogenic sources

Total Atmospheric Deposition to the Great Lakes Basin (ug/yr-person)

